



# **ESP CASE DETECTION ALGORITHM**

## **Human Immunodeficiency Virus (HIV)**

**Document Version 3.1**

**Prepared by the Department of Population Medicine at Harvard Medical School and Harvard Pilgrim Health Care Institute on behalf of the Massachusetts Department of Public Health.**

[esphealth@harvardpilgrim.org](mailto:esphealth@harvardpilgrim.org)

**January 8, 2020**



**Modification History**

Version	Date	Modification	By
3.1	1/8/2020	Added new HIV medications	DPM
3.0	3/1/2019	Added new provider reporting fields	DPM
2.5.1	1/24/2018	Added new HIV medication	DPM
2.5	9/14/2018	Added additional OI diagnosis codes	MDPH/DPM
2.4	12/8/2017	<ul style="list-style-type: none"> <li>Removed site-specific language</li> <li>Indicated which variables are not yet being reported via ESP.</li> <li>Added reporting time frames for lab results and medications.</li> </ul>	DPM
2.32	10/11/2017	Added ESP Logo and MDPH branding, formatting	DPM
2.31	10/4/2017	Transferred to updated algorithm template	DPM
2.3	7/14/2017	Added Multispot and Geenius tests to case criteria	MDPH/DPM
2.2	3/15/2017	<ul style="list-style-type: none"> <li>Added modified criteria to revoke a case.</li> <li>Minor modifications to reportable diagnosis codes.</li> </ul>	DPM
2.1	10/20/2016	Removed criteria to revoke a case.	DPM
2.0	9/22/2016	<ul style="list-style-type: none"> <li>Added conditions to the case criteria.</li> <li>Increased RNA Viral Load threshold.</li> </ul> Increased timeframe in which 3 different antivirals can occur.	DPM
1.0	5/10/2016	Original circulated version.	DPM



## Section 1. Overview

The purpose of this document is to describe the criteria used to identify and report Human Immunodeficiency Virus (HIV) cases and their continuum of care from electronic medical records (EMR) using ESP and report them to the Massachusetts Department of Public Health (MDPH). In addition, ESP will provide follow-up reports for subsequent events relevant to the continuum of care (e.g., serial CD4 and viral load testing, HIV treatment, etc.).

---

## Section 2. Criteria used to identify cases using ESP data

### I. CASE TYPES

This algorithm was developed to identify both new and prevalent HIV cases. Once identified, additional information will be captured on each case on an ongoing basis to evaluate the continuum of care.

### II. TIME WINDOW

Once HIV is acquired, cases are considered to be active lifelong. Thus, there is no recurrence window for this disease.

### III. CASE CRITERIA

Classify patient as HIV positive if any of the following conditions are true:

- A. Positive Western Blot, Multispot, or Geenius test result
- B. Positive HIV Antigen/Antibody test AND positive HIV ELISA (any time window)
- C. HIV RNA Viral Load > 200 copies/mL.
- D. HIV Qualitative PCR
- E.  $\geq 2$  ICD codes for HIV and history of prescription for  $\geq 3$  HIV meds ever
- F. HIV on problem list and history of prescription for  $\geq 3$  HIV meds ever
- G. Concurrent prescriptions for 3 or more *different* antiretrovirals for at least 1 month
  1. If patient prescribed combo pill then count as 2 or 3 meds as appropriate
  2. Rationale for prescription criteria:
    - a. Need med-based detection to capture patients on treatment with negative viral loads (since they'd be classified as negative using lab criteria alone)
    - b. 3 meds/3 months criteria to exclude patients on treatment for hepatitis B and patients receiving pre- or post-exposure prophylaxis
  3. Two approaches to meeting this criteria are valid: (a) based on the frequency and number of pills dispensed or (b) two start dates for each of 3 or more antiretrovirals at least 30 days apart but no more than 400 days apart.
    - a. The start dates for any set of 3 drugs need not be the same. For example, if 2 drugs are started Jan 1, and the 3<sup>rd</sup> drug is started Jan 10, this would count as meeting the 3 or more meds criteria as of Jan 10.
    - b. The sets of prescriptions for 3 or more antiretrovirals need not be for the same combination of drugs.

\*See [Section 4](#) for codes used to define each component.



**I. CRITERIA TO REVOKE A CASE**

For patients meeting criteria (e), (f), or (g) above, a case should be revoked (i.e., classified as HIV negative) if he/she subsequently has a negative ELISA or negative Ag/Ab. If there are  $\geq 2$  ELISA or Ag/Ab results on the same day, then only revoke if all tests are negative. Note, however, that these patients are eligible to become cases again if they subsequently meet one of the above criteria.

**Section 3. Specifications for reporting diseases/conditions to the Massachusetts Department of Public Health**

**I. INITIAL CASE REPORTING CRITERIA**

All cases will be reported to MDPH upon initial detection.

**II. CASE REPORT UPDATE CRITERIA**

A repeat report will be sent to MDPH whenever any of the following occur:

1. New CD4 test result
2. New viral load test result
3. Prescription for HIV med (new meds, not a renewal)
4. Encounter with diagnosis code for HIV
5. Encounter with a diagnosis code for an opportunistic infection

**III. DATA TO INCLUDE IN INITIAL REPORTS TO MDPH**

**A. Demographic**

Name	Last, first, middle
Date of birth	yyyy/mm/dd
Social security number	Last 4 digits
Gender	Male / Female
Race	American Indian / Asian / Black / White / Other / Unknown
Ethnicity	Hispanic / Non-Hispanic / Unknown
Address	Line 1, line 2, city, state, zip, country
Phone	xxx-xxx-xxxx
Language spoken	
Medical record number	
PCP	Name, office address, phone number, email
Country of birth	
Housing status	
Insurance status	

**B. Lab Ordering Facility Information**

Facility name	
---------------	--



Facility address	Line 1, line 2, city, state, zip
Facility contact person	
Contact person email	
Contact person phone	xxx-xxx-xxxx
Facility phone	xxx-xxx-xxxx
Facility NPI	

**C. Encounter Data**

See [Table 5](#) below for the NA codes to send provider fields in HL7 messages

Lab test ordering provider	Name and NPI
Prescribing provider	Name and NPI
Primary care provider	Name and NPI
Managing treatment provider	Name and NPI
Treatment encounter facility name	
Treatment encounter facility address	Line 1, line 2, city, state, zip
Treatment encounter facility NPI	
HIV visit date	Date of most recent encounter with an HIV code and all encounters following case identification

**D. Laboratory Results**

Report results within 30 days prior to the day on which case established and any time after the case date.

***Include test dates.***

HIV Antigen/Antibody Screen	
HIV ELISA Test (Antibody Screen)	
Western Blot	
Multispot	
Geenius	
HIV Quantitative RNA Viral Load	Most recent result
HIV Qualitative PCR	
CD4 Count	Most recent result

**E. Pregnancy**

Pregnancy flag active	Yes or No
Expected Date of Delivery (EDD)	

**F. Medications**

Report all medications from the specified ARV and OI medications lists given with 30 days prior to the case date and any time after the case date.

Treatment given	Yes or No
Current prescription for HIV medication	Text string including medication name, dose, frequency, and duration (calculated using start and end date). <a href="#">See Table 3</a> for specific medications.



Currently prescribed prophylactic medications	Text string including medication name, dose, frequency, and duration (calculated using start and end date). See <a href="#">Table 4</a> for specific medications.  <b>Note:</b> <i>OI medications will not be reported upon initial implementation. Modifications will need to be made in ESP to send these medications separately from the above.</i>
Treatment date	Date on which each prescription was ordered.

### G. Opportunistic Infections

If a diagnosis code for an opportunistic infection (see Table 2, below) are present within 30 days prior to case establishment, then report the following:

Infection name	<a href="#">See Table 2</a>
Date of last encounter with diagnosis code for opportunistic infection	

### H. Variables requiring further exploration

Several variables of interest for HIV still need to be explored with clinical providers. The following items are still under discussion, but will not hold up initial implementation.

Variable	Question in MAVEN	Status
Income level	“Income level as a percentage of poverty”	In the ESP model, but current reporting providers do not have this in their ESP data. Need to work with providers to understand if/how this information is captured.
Insurance type	“Type of health insurance?”	
Housing status	“Official Residence Type”	
Country of origin	“Country of birth”	
Sex risk	“HIV related risk”	Needs to be added to the ESP model and added by all providers.
Transgender status	“Gender”	Current reporting providers need to add “Transgender” as an option for gender in the ESP model.
Others	<ul style="list-style-type: none"> <li>• “Did the case attend their last scheduled HIV appointment?”</li> <li>• “Was resistance testing performed prior to initiating ARV?”</li> <li>• “Was the patient provided with risk reduction counseling?”</li> <li>• “Was patient referred to health department for partner services?”</li> <li>• “Date of HIV Care Coordination Event” (separate from “HIV clinical visit date”)</li> </ul>	Need to work with providers to understand if/how this information is captured. These variables would need to be added to the ESP model.



**Section 4. Codes, laboratory tests, and medications used to identify criteria listed in Sections 1 and 2.**

**I. DIAGNOSES CODES**

**Table 1. Diagnosis Codes Used to Identify HIV**

Code Type	Code	Description
ICD-9-CM	042	HIV disease
ICD-9-CM	V08	Asymptomatic HIV infection status
ICD-10-CM	B20-B24	Human immunodeficiency virus (HIV) disease
ICD-10-CM	Z21	Asymptomatic HIV infection status
ICD-10-CM	B97.35	Human immunodeficiency virus, type 2 [HIV 2] as the cause of diseases classified elsewhere
ICD-10-CM	O98.7	HIV complicating pregnancy or childbirth

**Table 2. Diagnosis Codes Used to define Opportunistic Infection**

Code Type	Code	Description
ICD-9-CM	136.3	PCP pneumonia
ICD-10-CM	B59	PCP pneumonia
ICD-9-CM	130.*	Toxoplasmosis
ICD-10-CM	B58, B58.2, B58.9	Toxoplasmosis
ICD-9-CM	117.5, 321.0	Cryptococcus and Cryptococcal meningitis
ICD-10-CM	B45, B45.0, B45.1, B45.2, B45.3, B45.7, B45.8, B45.9	Cryptococcus and Cryptococcal meningitis
ICD-9-CM	007.4	Cryptosporidiosis
ICD-10-CM	A07.2	Cryptosporidiosis
ICD-9-CM	010-018	Tuberculosis
ICD-10-CM	A15-A19.9	Tuberculosis
ICD-9-CM	112.0	Candidiasis of mouth
ICD-9-CM	112.4	Candidiasis of lung
ICD-9-CM	112.5	Disseminated candidiasis
ICD-9-CM	112.84	Candidal esophagitis
ICD-10-CM	B37.0	Candidal stomatitis
ICD-10-CM	B37.1	Pulmonary candidiasis
ICD-10-CM	B37.81	Candidal esophagitis
ICD-10-CM	B37.83	Candidal cheilitis
ICD-9-CM	031.2	Disseminated MAC
ICD-10-CM	A31.2	Disseminated MAC
ICD-10-CM	A31, A31.0, A31.1, A31.8, A31.9	Mycobacterium infection
ICD-9-CM	046.3	Progressive multifocal leukencephalopathy (JC Virus)
ICD-10-CM	A81.2	Progressive multifocal leukencephalopathy (JC Virus)
ICD-10-CM	D06, D06.0, D06.1, D06.7, D06.9	Invasive cervical carcinoma



ICD-10-CM	B38, B38.0, B38.1, B38.2, B38.3, B38.4, B38.7, B38.8, B38.81, B38.89, B38.9	Coccidioidomycosis
ICD-10-CM	B25.0, B25.2, B25.8, B25.9	Cytomegalovirus disease
ICD-10-CM	G93.40, G93.49	Encephalopathy
ICD-10-CM	B00, B00.0, B00.1, B00.2, B00.3, B00.4, B00.5, B00.50, B00.51, B00.52, B00.53, B00.59, B00.7, B00.8, B00.81, B00.82, B00.89, B00.9	Herpesviral infection
ICD-10-CM	B39, B39.0, B39.1, B39.2, B39.3, B39.4, B39.5, B39.9	Histoplasmosis
ICD-10-CM	A07.3	Isosporiasis
ICD-10-CM	C46, C46.0, C46.1, C46.2, C46.3, C46.4, C46.5, C46.50, C46.51, C46.52, C46.7, C46.9	Kaposi sarcoma
ICD-10-CM	J84.2	Lymphoid interstitial pneumonia
ICD-10-CM	C83.70, C83.71, C83.72, C83.73, C83.74, C83.75, C83.76, C83.77, C83.78, C83.79	Burkitt lymphoma
ICD-10-CM	C83.30, C83.39, C83.80, C83.89	Diffuse large B-cell lymphoma
ICD-10-CM	J09.X1, J10.0, J11.0, J12, J12.0, J12.1, J12.2, J12.3, J12.8, J12.81, J12.89, J12.9, J13, J16, J16.0, J16.8, J17, J18, J18.0, J18.1, J18.2, J18.8, J18.9	Recurrent pneumonia
ICD-10-CM	A02.1	Salmonella sepsis
ICD-10-CM	A81.2	Progressive multifocal leukoencephalopathy
ICD-10-CM	R64, M62.5, M62.50, M62.58, M62.59	Wasting syndrome

## II. MEDICATIONS

**Table 3. HIV Medications**

Drug Class	Generic Name	Brand Name
NRTI	Zidovudine (AZT)	Retrovir
	Didanosine (DDI)	Videx
	Stavudine (D4T)	Zerit
	Lamivudine (3TC)	Epivir
	Emtricitabine (FTC)	Emtriva
	Tenofovir (TDF)	Viread
	Abacavir (ABC)	Ziagen
	Tenofovir + Emtricitabine	Truvada
	Zidovudine + Lamivudine	Combivir



	Abacavir + Lamivudine	Epzicom
	Abacavir + Lamivudine + Zidovudine	Trizivir
	Emtricitabine + Tenofovir alafenamide	Descovy
NNRTIs	Efavirenz	Sustiva
	Nivirapine	Viramune
	Rilpivirine	Eduvant
	Etravirine	Intelence
	Delavirdine	Rescriptor
Integrase Inhibitors	Raltegravir	Isentress
	Dolutegravir	Tivicay
	Elvitegravir	
Fusion Inhibitors	Enfuvirtide	Fuzeon
	Maraviroc	Selzentry
Protease Inhibitors	Tipranavir	Aptivus
	Ritonavir	Norvir
	Indinavir	Crixivan
	Darunavir	Prezista
	Saquinavir	Invirase
	Atazanavir	Reyataz
	Nelfinavir	Viracept
	Fosamprenavir	Lexiva
	Lopinavir + Ritonavir	Kaletra
Single Tablet Combos	Efavirenz + Tenofovir + Emtricitabine	Atripla
	Rilpivirine + Tenofovir + Emtricitabine	Complera
	Dolutegravir + Abacavir + Lamivudine	Triumeq
	Elvitegravir + Cobicistat + Tenofovir + Emtricitabine	Stribild
	Dolutegravir + Cobicistat	Prezcobix
	Bictegravir + Emtricitabine + Tenofovir alafenamide	Biktarvy
	Elvitegravir + Cobicistat + Emtricitabine + Tenofovir alafenamide	Genvoya
	Emtricitabine + Rilpivirine + Tenofovir alafenamide	Odefsey

**Table 4. Medications for Primary Prevention of Opportunistic Infections**

Infection	Generic Name	Brand Name
Pneumocystis	TMP-SMX	Bactrim
	Dapsone	None (?)
	Pyrimethamine	Daraprim
	Atovaquone	Mepron
	Pentamidine	Nebupent, Pentam 300
Toxoplasmosis	TMP-SMX	
	Dapsone	
	Pyrimethamine	
MAC	Azithromycin	Zithromax, Zmax



### III. LABORATORY TESTS LOINCS

**Table 4. Laboratory Test LOINC Mapping**

Component Name	LOINC	LOINC Name	Positive Result
Antigen/antibody combination immunoassay	56888-1	HIV 1+2 Ab+HIV1 p24 Ag : ACnc : Pt : Ser : Ord : EIA	
Antibody Screen, HIV1	29327-4	HIV1 Ab : ACnc : Pt : Body fld : QI : Ord	
Antibody Screen, HIV2	30361-0	HIV2 Ab [Presence] : ACnc: Pt : Ser : QI : Ord : EIA	
Antibody Screen, HIV1+2	43010-8	HIV 1+2 Ab [Presence]: ACnc : Pt : XXX : Ord	
CD4 (#/volume)	32515-9	Deprecated CD4 cells [# /volume]: NCnc : Pt: XXX : Qn	
CD4 (/100cells)	32516-7	Deprecated CD4 cells [/100 Cells] : NFr : Pt: XXX : Qn	
Qualitative Viral Load	5018-7	HIV 1 RNA [Presence] : ACnc : Pt : XXX : Ord : Probe.amp.tar	
Quantitative Viral Load	25836-8	HIV 1 RNA [# /volume] (viral load) : NCnc : Pt : XXX : Probe.amp.tar	
Western Blot, HIV1+2	43185-8	HIV 1 & 2 Ab band pattern [interpretation] : Imp : Pt : Ser : Nom : IB	
Western Blot, HIV1	34592-6	HIV 1 Ab [Presence] : ACnc : Pt : Body fld : Ord : IB	
Western Blot, HIV2	5225-8	HIV 2 Ab [Presence] : ACnc : Pt: Ser : Ord : IB	
Differentiating Multispot	69668-2	HIV 1 and 2 Ab [Identifier] in Serum or Plasma by Rapid immunoassay	HIV Type 1 HIV Type 2 Positive
Differentiating Geenius	80203-3	HIV 1 and 2 Ab in Serum/Plasma/Blood by Rapid immunoassay	HIV Type 1 HIV Type 2 HIV Type 1 Indeterminate HIV Type 2 Indeterminate HIV Type 2 with HIV 1 Crossreactivity Positive (Untypable)

### IV. NA CODES

**Table 5. NA Codes for provider reporting fields**

NA code	Description
NA-1746	Prescribing provider name
NA-1747	Prescribing provider NPI
NA-1748	Treatment encounter facility name
NA-1749	Treatment encounter facility address
NA-1750	Treatment encounter facility city
NA-1751	Treatment encounter facility state
NA-1752	Treatment encounter facility NPI
NA-1753	Primary care provider name
NA-1754	Primary care provider NPI
NA-1755	Ordering provider name
NA-1756	Ordering provider NPI
NA-1757	Lab ordering facility name
NA-1758	Lab ordering facility address
NA-1759	Lab ordering facility city
NA-1760	Lab ordering facility state



NA-1761	Lab ordering facility NPI
NA-1762	Managing treatment provider name
NA-1763	Managing treatment provider NPI

## V. CODE MAINTENANCE STRATEGY

Search all incoming new labs for strings that might indicate relevance to HIV including:

- HIV
- Immunodef
- CD4